

Lawn Care Calendar

Bermudagrass

Aaron Patton
Assistant Professor -
Turfgrass Specialist

John Boyd
Professor -
Weed Scientist

These suggested maintenance practices will help you care for your lawn throughout the year. Because every site is different due to variations in location, terrain, soil type, condition of lawn, previous lawn care and other factors, adjust these practices and dates to suit your home lawn.

Bermudagrass (*Cynodon* spp.) is the most commonly used lawn grass in Arkansas. It will grow on a wide range of soil types as long as there is adequate drainage and plenty of sunlight. Bermudagrass is not a shade-tolerant turfgrass. Full sun is required for it to thrive. Other attractive features include rapid recovery from traffic damage and good drought tolerance. Established bermudagrass will turn brown during extended dry periods but recover after the first significant rainfall. Durability and the ability to recover quickly make it the first choice for high-traffic areas.

The quick growth rate of bermudagrass compared to slower-growing grasses like zoysiagrass and centipedegrass makes bermudagrass the most affordable turfgrass to purchase as sod. Rapid growth means frequent mowing during June, July and August. Mowing frequency is also tied to nitrogen fertilization and rainfall or irrigation. Because bermudagrass is a fast grower that produces rhizomes and stolons, it readily invades ornamental beds, gardens and requires frequent edging along walks and driveways. The aggressive properties that make it a desirable turfgrass also make it a major weed.

The hybrid bermudagrasses (*Cynodon dactylon* × *C. transvaalensis*), which include ‘Tifway’ (Tifton 419), ‘TifSport’, ‘Patriot’

and many others, are generally finer textured than common bermudagrass (*Cynodon dactylon*) and must be started from sod, sprigs or plugs. For more information on locating bermudagrass cultivars, see the *Arkansas Sod Source Directory*, FSA6136.

Common bermudagrass is typically seeded. After years of nothing but ‘Arizona Common’ bermudagrass on the market, the quality of seeded bermudagrasses has improved dramatically since 2000. New seeded varieties such as ‘Riviera’ and ‘Yukon’ (developed at Stillwater, Oklahoma) are a good choice for lawns in Arkansas. They are attractive, cold-tolerant grasses that are well adapted in all parts of the state. ‘Princess-77’ (from New Mexico) is a seeded, fine-textured variety that approaches the quality of ‘Tifway’, but lacks sufficient cold tolerance for use in north Arkansas. Many other cultivars are also available. For more information on seeding a lawn, see *Seeding a Lawn in Arkansas*, FSA2113.

Tifway is a hybrid released in 1960 from the USDA research station in Tifton, Georgia. Tifway is available from many sod farms in Arkansas. While not ideal, Tifway is a better choice than ‘Tifgreen’ (Tifton 328) for lawns in Arkansas. The biggest problem that homeowners have with hybrid bermudagrasses is the inability to mow them correctly. Hybrid bermudagrasses look best when mowed three times per week at 0.5 to 1.5 inch with a reel mower. Because this isn’t practical for most homeowners, a seeded bermudagrass is often a better choice.

*Arkansas Is
Our Campus*

Visit our web site at:
<http://www.uaex.edu>

Bermudagrass maintenance calendar.

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Watering†				●	●	●	●	●	●	●		
Mowing				●	●	●	●	●	●	●		
Fertilization				●	●	●	●	●	●			
Liming		●	●	●	●				●	●	●	●
Aeration				●	●	●	●	●	●			
Dethatching				●	●	●	●	●				
Seeding		●	●	●	●	●						
Sodding		●	●	●	●	●	●	●	●	●	●	●
<hr/>												
Weed control												
Preemergence-crabgrass	●	●	●	●								
Postemergence-broadleaf			●	●	●	●	●		●	●	●	
Postemergence-grasses/sedges				●	●	●	●	●				
Postemergence-winter annuals											●	●

† Bullets represent the optimum time period to perform various maintenance practices to your bermudagrass lawn. The optimum lawn maintenance period may be started earlier or extended based on variations in annual weather conditions and/or location in Arkansas. Dark bullets represent the best months for each practice, and lightly shaded bullets represent possible months.

	March Through May
Mowing	<p>Before bermudagrass begins to grow in the spring, you may mow the turf slightly shorter than normal to remove dead leaf blades and other debris. This practice will reduce shading of the emerging plants and also serve to warm soil temperatures more quickly in the spring. Begin mowing regularly when the grass turns green in the spring and reaches the desired mowing height. Bermudagrass should be mowed at 0.5 to 2.5 inches. Hybrid bermudagrass cultivars should be mowed at heights from 0.5 to 1.5 inches. Common bermudagrass should be mowed at 1.5 to 2.5 inches. Higher mowing heights may be needed on uneven lawns to avoid scalping. Mow often to avoid removing more than one-third of the leaf blade. It isn't necessary to collect clippings unless they remain as clumps on the lawn surface. Mowing more frequently to avoid clipping buildup is more efficient than emptying the collection bag. For more information about mowing, see <i>Mowing Your Lawn</i>, FSA6023.</p>
Fertilizing	<p>Apply 0.5 to 1.0 pound of nitrogen per thousand square feet approximately three weeks after the grass turns green in late May. Submit a soil sample to determine phosphorus and potassium requirements, if you haven't already (contact your county Extension office). Apply lime if suggested. For more information about fertilization, see <i>Fertilizing Your Lawn</i>, FSA2114. For more information about soil pH in lawns, see <i>Liming Your Lawn</i>, FSA6134.</p> <p>To determine the amount of fertilizer product required to apply 1.0 pound of nitrogen per thousand square feet, divide 1.0 by the first number (%) in the fertilizer ratio. For example, for a 20-5-5 fertilizer (containing 20% nitrogen), divide 1.0 by 0.20 (NOTE: 20% = 0.20). The result is 5 pounds of product per thousand square feet. For more information on calculating the amount of fertilizer you need to apply, see <i>Fertilizing Your Lawn</i>, FSA2114.</p>
Watering	<p>Irrigation is seldom needed on bermudagrass during the spring except for newly sodded areas or if dry, hot, windy conditions occur for an extended period. A dark, bluish gray color, foot-printing and wilted, folded or curled leaves indicate it is time to water. Proper irrigation may prevent or reduce pest problems and environmental stress later in the summer.</p>

March Through May (cont.)	
Weed Control	Apply preemergence herbicides to control crabgrass in late February or early March. Apply postemergence herbicides in May as needed to control summer annual and perennial broadleaf weeds such as knotweed, spurge and lespedeza. Products containing two or three broadleaf herbicides usually control several different broadleaf weeds in a lawn more effectively. Be sure the product is labeled for use on bermudagrass. Apply postemergence herbicides only when weeds are present. Applying broadleaf herbicides three weeks after the lawn becomes green to avoid damaging the bermudagrass. See FSA2109, <i>Home Lawn Weed Control</i> , for more weed control information.
Disease Control	Small (1- to 3-inch) circular, shaded, straw-colored spots in your lawn appearing several weeks after greenup are likely an indicator of the disease dollar spot. Larger dead spots (1 to 3 feet) in diameter appearing during spring greenup and for several weeks following likely indicate the presence of spring dead spot. For more information about these diseases, see FSA7541, <i>Dollar Spot of Turfgrass in the Home Landscape</i> , and FSA7551, <i>Bermudagrass Spring Dead Spot</i> .
Insect Control	Check for insect pests and treat if necessary.
Renovation	Replant large bare areas using sod or plugs planted on 6- or 12-inch centers. Applying a preemergence herbicide that does not interfere with root growth after plugging helps prevent weed encroachment. Common bermudagrass can be seeded at 0.5 to 1 pound per thousand square feet in lawns not treated with a preemergence herbicide within the previous two months.
June Through August	
Mowing	Bermudagrass should be mowed every 5 to 7 days and less often when the lawn is drought stressed.
Fertilizing	Apply 0.5 to 1.0 pound of nitrogen per thousand square feet every 4 to 6 weeks using the March through May fertilizing guidelines. The interval between fertilizer applications may be increased by applying a slow-release fertilizer.
Watering	Water early in the morning to wet the soil to a depth of 4 to 6 inches. Probe with a screwdriver to determine moisture depth. Bermudagrass needs a weekly application of 1 to 1.25 inches of water to retain its color during summer. It needs even less to survive and can go several weeks without supplemental irrigation. On sandy soils, it requires more frequent watering: for example, 0.5 inch of water every third day. It is often necessary to irrigate an area for three to five hours to apply 1 inch of water with most homeowner irrigation systems. (It takes 620 gallons of water to apply 1 inch of water per thousand square feet.) Because clay soils accept water slowly, irrigate these areas until runoff occurs; wait one-half hour until the water has been absorbed, and then continue irrigating until the desired depth or amount is obtained. A dark, bluish gray color, foot-printing and wilted, folded or curled leaves indicate that it is time to water.
Cultivation	Thatch needs to be removed every two to three years through core aeration or dethatching. Cultivation during the early summer is preferred since moisture is usually not limiting and growth is optimum for recovery. For more information about thatch, see <i>Thatch Prevention and Control</i> , FSA6139.
Insect Control	Check for insect pests and treat if necessary.
Weed Control	Apply postemergence herbicides as needed to control summer annual and perennial broadleaf weeds such as knotweed, spurge and lespedeza. Crabgrass, dallisgrass, annual sedges and sandbur can be controlled with MSMA. Two applications 5 to 7 days apart are required for effective annual sedge and crabgrass control. Dallisgrass requires five applications of MSMA 5 to 7 days apart. Apply herbicides only when weeds are present, the grass is actively growing and the lawn is not suffering from drought stress.
Renovation	Replant large bare areas using sod or plugs planted on 6- or 12-inch centers. Applying a preemergence herbicide that does not interfere with root growth after plugging helps prevent weed encroachment. Common bermudagrass can be seeded at 0.5 to 1 pound per thousand square feet. Seeding in spring or early summer will enhance the winter hardiness of bermudagrass seedlings.

September Through November	
Mowing	Bermudagrass should be mowed every 5 to 7 days and less often when the lawn is drought stressed.
Fertilizing	Apply no more than 0.5 pound of nitrogen per thousand square feet in September, four to six weeks before the first expected frost. Potassium can be applied if soil tests indicate a need. If potassium is needed, use a low-nitrogen, high-potassium fertilizer such as a 10-0-40 or supplement a nitrogen fertilizer source with 1 pound of potash using 1.6 pounds of muriate of potash (0-0-60) per 1,000 square feet. Several manufacturers offer winterizing fertilizers with various combinations of nutrients. When using these products, try to find one that approximates at 4-1-6 ratio of nitrogen-phosphorous-potassium and contains iron, which will extend color into fall. Apply lime during these months if recommended by your soil test.
Watering	Follow the March through May irrigation guidelines. Dormant bermudagrass may still need to be watered periodically when dry, windy conditions occur for an extended period. Additionally, newly planted sod should be watered during this period to prevent desiccation.
Disease Control	Some diseases such as spring dead spot are active in the fall despite the fact that no symptoms are visible. If your lawn has a history of spring dead spot, fall applications of fungicides may be beneficial. For more information about this disease, see FSA7551, <i>Bermudagrass Spring Dead Spot</i> .
Weed Control	Apply preemergence or postemergence herbicides as needed to control winter annual and perennial broadleaf weeds such as chickweed and henbit. Preemergence herbicides do not control existing perennial weeds. Apply postemergence herbicides only when weeds are present. Do not apply herbicides designed to control annual bluegrass if the lawn is to be overseeded with ryegrass.
December Through February	
Mowing	Mow bermudagrass overseeded with perennial or annual ryegrass as often as needed not to remove more than one-third of the leaf blade. Recycle nutrients by not collecting the clippings unless they accumulate heavily on the surface. Dormant bermudagrass that has not been overseeded does not need mowing.
Fertilizing	Do not fertilize bermudagrass that has not been overseeded. For overseeded bermudagrass, apply ½ pound of nitrogen per thousand square feet in December and February. In the absence of a soil test, use a complete (N-P-K) turf-grade fertilizer with a 3-1-2 or 4-1-2 ratio (for example, 12-4-8 or 24-6-12). Submit soil samples for analysis every 2-3 years to determine your lawn's nutrient requirements. Be sure to specify your lawn species.
Watering	Newly planted sod should be watered during this period to prevent desiccation. Watering is particularly important for lawns that have been overseeded.
Weed Control	Apply broadleaf herbicides as needed to control weed such as spurweed, chickweed, henbit and hop clover. Apply postemergence herbicides only when weeds are present. Selective herbicides like atrazine and simazine can be applied in November or December in non-overseeded lawns to control annual bluegrass and several winter annual broadleaf weeds such as henbit.

Additional Information

Additional fact sheets available at <http://www.uaex.edu>

Additional information about turfgrass management available at <http://turf.uark.edu>

Printed by University of Arkansas Cooperative Extension Service Printing Services.

DR. AARON PATTON is assistant professor - turfgrass specialist with the University of Arkansas Division of Agriculture, Cooperative Extension Service, in Fayetteville. **DR. JOHN BOYD** is professor - weed scientist with the University of Arkansas Division of Agriculture, Cooperative Extension Service, in Little Rock.

FSA6121-PD-11-08RV

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The Arkansas Cooperative Extension Service offers its programs to all eligible persons regardless of race, color, national origin, religion, gender, age, disability, marital or veteran status, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.